

# Generalized Harmonic Evolutions of Binary Black Hole Spacetimes

Lee Lindblom  
Caltech

Collaborators: Michael Boyle, Duncan Brown, Gregory Cook,  
Larry Kidder, Geoffrey Lovelace, Abdul Mroue, Robert Owen,  
Harald Pfeiffer, Oliver Rinne, Mark Scheel, Saul Teukolsky

University of Wisconsin at Milwaukee — 4 May 2007

# Generalized Harmonic Evolutions of Binary Black Hole Spacetimes

Lee Lindblom  
Caltech

Collaborators: Michael Boyle, Duncan Brown, Gregory Cook,  
Larry Kidder, Geoffrey Lovelace, Abdul Mroue, Robert Owen,  
Harald Pfeiffer, Oliver Rinne, Mark Scheel, Saul Teukolsky

University of Wisconsin at Milwaukee — 4 May 2007

- Generalize Harmonic (GH) gauge conditions.

# Generalized Harmonic Evolutions of Binary Black Hole Spacetimes

Lee Lindblom  
Caltech

Collaborators: Michael Boyle, Duncan Brown, Gregory Cook, Larry Kidder, Geoffrey Lovelace, Abdul Mroue, Robert Owen, Harald Pfeiffer, Oliver Rinne, Mark Scheel, Saul Teukolsky

University of Wisconsin at Milwaukee — 4 May 2007

- Generalize Harmonic (GH) gauge conditions.
- Constraint damping in the GH system.

# Generalized Harmonic Evolutions of Binary Black Hole Spacetimes

Lee Lindblom  
Caltech

Collaborators: Michael Boyle, Duncan Brown, Gregory Cook, Larry Kidder, Geoffrey Lovelace, Abdul Mroue, Robert Owen, Harald Pfeiffer, Oliver Rinne, Mark Scheel, Saul Teukolsky

University of Wisconsin at Milwaukee — 4 May 2007

- Generalize Harmonic (GH) gauge conditions.
- Constraint damping in the GH system.
- Moving Black Holes.

# Generalized Harmonic Evolutions of Binary Black Hole Spacetimes

Lee Lindblom  
Caltech


Collaborators: Michael Boyle, Duncan Brown, Gregory Cook,  
Larry Kidder, Geoffrey Lovelace, Abdul Mroue, Robert Owen,  
Harald Pfeiffer, Oliver Rinne, Mark Scheel, Saul Teukolsky

University of Wisconsin at Milwaukee — 4 May 2007

- Generalize Harmonic (GH) gauge conditions.
- Constraint damping in the GH system.
- Moving Black Holes.
- Binary Black Hole Evolutions.

# Traditional ADM Gauge Conditions

- Construct a foliation of spacetime by spatial slices.
- Choose a time function with  $t = \text{const.}$  on these slices.
- Choose spatial coordinates,  $x^k$ , on each slice.
- Decompose the 4-metric  $\psi_{ab}$  into its 3+1 parts:  
$$ds^2 = \psi_{ab} dx^a dx^b = -N^2 dt^2 + g_{ij} (dx^i + N^i dt)(dx^j + N^j dt).$$
- The lapse  $N$  and shift  $N^i$  measure how coordinates are laid out on spacetime:  
$$\begin{aligned} \vec{n} = \partial_\tau &= \frac{\partial t}{\partial \tau} \partial_t + \frac{\partial x^k}{\partial \tau} \partial_k, \\ &= \frac{1}{N} \partial_t - \frac{N^k}{N} \partial_k. \end{aligned}$$
- Spacetime coordinates are determined in the traditional ADM method by specifying the lapse  $N$  and shift  $N^i$ .



# Generalized Harmonic Gauge Conditions

- An alternate way to specify the coordinates is through the generalized harmonic gauge source function  $H^a$ :
- Let  $H^a$  denote the function obtained by the action of the scalar wave operator on the coordinates  $x^a$ :

$$H^a \equiv \nabla^c \nabla_c x^a = \psi^{bc} (\partial_b \partial_c x^a - \Gamma_{bc}^e \partial_e x^a) = -\Gamma^a,$$

where  $\Gamma^a = \psi^{bc} \Gamma_{bc}^a$  and  $\psi_{ab}$  is the 4-metric.

# Generalized Harmonic Gauge Conditions

- An alternate way to specify the coordinates is through the generalized harmonic gauge source function  $H^a$ :
- Let  $H^a$  denote the function obtained by the action of the scalar wave operator on the coordinates  $x^a$ :

$$H^a \equiv \nabla^c \nabla_c x^a = \psi^{bc} (\partial_b \partial_c x^a - \Gamma_{bc}^e \partial_e x^a) = -\Gamma^a,$$

where  $\Gamma^a = \psi^{bc} \Gamma^a_{bc}$  and  $\psi_{ab}$  is the 4-metric.

- Specifying coordinates by the *generalized harmonic* (GH) method can be accomplished by choosing a gauge-source function  $H_a(x, \psi) = \psi_{ab} H^b$ , and requiring that  $H_a(x, \psi) = -\Gamma_a = -\Gamma_{abc} \psi^{bc}$ .



## Important Properties of the GH Method

- The Einstein equations are manifestly hyperbolic when coordinates are specified using a GH gauge function:

$$R_{ab} = -\frac{1}{2}\psi^{cd}\partial_c\partial_d\psi_{ab} + \nabla_{(a}\Gamma_{b)} + F_{ab}(\psi, \partial\psi),$$

where  $\psi_{ab}$  is the 4-metric, and  $\Gamma_a = \psi^{bc}\Gamma_{abc}$ . The vacuum Einstein equation,  $R_{ab} = 0$ , has the same principal part as the scalar wave equation when  $H_a(\mathbf{x}, \psi) = -\Gamma_a$  is imposed.

# Important Properties of the GH Method

- The Einstein equations are manifestly hyperbolic when coordinates are specified using a GH gauge function:

$$R_{ab} = -\frac{1}{2}\psi^{cd}\partial_c\partial_d\psi_{ab} + \nabla_{(a}\Gamma_{b)} + F_{ab}(\psi, \partial\psi),$$

where  $\psi_{ab}$  is the 4-metric, and  $\Gamma_a = \psi^{bc}\Gamma_{abc}$ . The vacuum Einstein equation,  $R_{ab} = 0$ , has the same principal part as the scalar wave equation when  $H_a(x, \psi) = -\Gamma_a$  is imposed.

- Imposing coordinates using a GH gauge function profoundly changes the constraints. The GH constraint,  $\mathcal{C}_a = 0$ , where

$$\mathcal{C}_a = H_a + \Gamma_a,$$

depends only on first derivatives of the metric. The standard Hamiltonian and momentum constraints,  $\mathcal{M}_a = 0$ , are determined by the derivatives of the gauge constraint  $\mathcal{C}_a$ :

$$\mathcal{M}_a \equiv G_{ab}n^b = \left[ \nabla_{(a}\mathcal{C}_{b)} - \frac{1}{2}\psi_{ab}\nabla^c\mathcal{C}_c \right] n^b.$$

# Constraint Damping Generalized Harmonic System

- Pretorius (based on a suggestion from Gundlach, et al.) modified the GH system by adding terms proportional to the gauge constraints:

$$0 = R_{ab} - \nabla_{(a} \mathcal{C}_{b)} + \gamma_0 \left[ n_{(a} \mathcal{C}_{b)} - \frac{1}{2} \psi_{ab} n^c \mathcal{C}_c \right],$$

where  $n^a$  is a unit timelike vector field. Since  $\mathcal{C}_a = H_a + \Gamma_a$  depends only on first derivatives of the metric, these additional terms do not change the hyperbolic structure of the system.

# Constraint Damping Generalized Harmonic System

- Pretorius (based on a suggestion from Gundlach, et al.) modified the GH system by adding terms proportional to the gauge constraints:

$$0 = R_{ab} - \nabla_{(a} \mathcal{C}_{b)} + \gamma_0 \left[ n_{(a} \mathcal{C}_{b)} - \frac{1}{2} \psi_{ab} n^c \mathcal{C}_c \right],$$

where  $n^a$  is a unit timelike vector field. Since  $\mathcal{C}_a = H_a + \Gamma_a$  depends only on first derivatives of the metric, these additional terms do not change the hyperbolic structure of the system.



- Evolution of the constraints  $\mathcal{C}_a$  follow from the Bianchi identities:

$$0 = \nabla^c \nabla_c \mathcal{C}_a - 2\gamma_0 \nabla^c [n_{(c} \mathcal{C}_{a)}] + \mathcal{C}^c \nabla_{(c} \mathcal{C}_{a)} - \frac{1}{2} \gamma_0 n_a \mathcal{C}^c \mathcal{C}_c.$$

This is a damped wave equation for  $\mathcal{C}_a$ , that drives all small short-wavelength constraint violations toward zero as the system evolves (for  $\gamma_0 > 0$ ).


# Numerical Tests of the New GH System

- 3D numerical evolutions of static black-hole spacetimes illustrate the constraint damping properties of our GH evolution system.
- These evolutions are stable and convergent when  $\gamma_0 = \gamma_2 = 1$ .




# Moving Black Holes in a Spectral Code

- Spectral: Excision boundary is a smooth analytic surface.




# Moving Black Holes in a Spectral Code

- Spectral: Excision boundary is a smooth analytic surface.
  - Cannot add/remove individual grid points.



# Moving Black Holes in a Spectral Code


- Spectral: Excision boundary is a smooth analytic surface.
  - Cannot add/remove individual grid points.
- Straightforward method: re-grid when holes move too far.






# Moving Black Holes in a Spectral Code

- Spectral: Excision boundary is a smooth analytic surface.
  - Cannot add/remove individual grid points.
- Straightforward method: re-grid when holes move too far.




# Moving Black Holes in a Spectral Code

- Spectral: Excision boundary is a smooth analytic surface.
  - Cannot add/remove individual grid points.
- Straightforward method: re-grid when holes move too far.




# Moving Black Holes in a Spectral Code

- Spectral: Excision boundary is a smooth analytic surface.
  - Cannot add/remove individual grid points.
- Straightforward method: re-grid when holes move too far.
- **Problems:**
  - Re-gridding/interpolation is expensive.




# Moving Black Holes in a Spectral Code

- Spectral: Excision boundary is a smooth analytic surface.
  - Cannot add/remove individual grid points.
- Straightforward method: re-grid when holes move too far.
- **Problems:**
  - Re-gridding/interpolation is expensive.
  - Difficult to get smooth extrapolation at trailing edge of horizon.





# Moving Black Holes in a Spectral Code

- Spectral: Excision boundary is a smooth analytic surface.
  - Cannot add/remove individual grid points.
- Straightforward method: re-grid when holes move too far.
- **Problems:**
  - Re-gridding/interpolation is expensive.
  - Difficult to get smooth extrapolation at trailing edge of horizon.
  - Causality trouble at leading edge of horizon.





# Moving Black Holes in a Spectral Code

- Spectral: Excision boundary is a smooth analytic surface.
  - Cannot add/remove individual grid points.
- Straightforward method: re-grid when holes move too far.
- **Problems:**
  - Re-gridding/interpolation is expensive.
  - Difficult to get smooth extrapolation at trailing edge of horizon.
  - Causality trouble at leading edge of horizon.



# Moving Black Holes in a Spectral Code

- Spectral: Excision boundary is a smooth analytic surface.
  - Cannot add/remove individual grid points.
- Straightforward method: re-grid when holes move too far.
- **Problems:**
  - Re-gridding/interpolation is expensive.
  - Difficult to get smooth extrapolation at trailing edge of horizon.
  - Causality trouble at leading edge of horizon.




# Moving Black Holes in a Spectral Code

- Spectral: Excision boundary is a smooth analytic surface.
  - Cannot add/remove individual grid points.
- Straightforward method: re-grid when holes move too far.
- **Problems:**
  - Re-gridding/interpolation is expensive.
  - Difficult to get smooth extrapolation at trailing edge of horizon.
  - Causality trouble at leading edge of horizon.

- **Solution:**

Choose coordinates that smoothly track the location of the black hole.


For a black hole binary this means using coordinates that rotate with respect to inertial frames at infinity.





# Evolving Black Holes in Rotating Frames

- Coordinates that rotate with respect to the inertial frames at infinity are needed to track the horizons of orbiting black holes.
- Evolutions of Schwarzschild in rotating coordinates are unstable.



- Evolutions shown use a computational domain that extends to  $r = 1000M$ .
- Angular velocity needed to track the horizons of an equal mass binary at merger is about  $\Omega \approx 0.2/M$ .
- Problem caused by asymptotic behavior of metric in rotating coordinates:  $\psi_{tt} \sim \rho^2 \Omega^2$ ,  $\psi_{ti} \sim \rho \Omega$ ,  $\psi_{ij} \sim 1$ .

## Dual-Coordinate-Frame Evolution Method

- Single-coordinate frame method uses the one set of coordinates,  $x^{\bar{a}} = \{\bar{t}, x^{\bar{i}}\}$ , to define field components,  $u^{\bar{\alpha}} = \{\psi_{\bar{a}\bar{b}}, \Pi_{\bar{a}\bar{b}}, \Phi_{\bar{i}\bar{a}\bar{b}}\}$ , and the same coordinates to determine these components by solving Einstein's equation for  $u^{\bar{\alpha}} = u^{\bar{\alpha}}(x^{\bar{a}})$ :

$$\partial_{\bar{i}} u^{\bar{\alpha}} + A^{\bar{k}\bar{\alpha}}_{\bar{\beta}} \partial_{\bar{k}} u^{\bar{\beta}} = F^{\bar{\alpha}}.$$

# Dual-Coordinate-Frame Evolution Method

- Single-coordinate frame method uses the one set of coordinates,  $x^{\bar{a}} = \{\bar{t}, x^{\bar{i}}\}$ , to define field components,  $u^{\bar{\alpha}} = \{\psi_{\bar{a}\bar{b}}, \Pi_{\bar{a}\bar{b}}, \Phi_{\bar{i}\bar{a}\bar{b}}\}$ , and the same coordinates to determine these components by solving Einstein's equation for  $u^{\bar{\alpha}} = u^{\bar{\alpha}}(x^{\bar{a}})$ :

$$\partial_{\bar{i}} u^{\bar{\alpha}} + A^{\bar{k}\bar{\alpha}}_{\bar{\beta}} \partial_{\bar{k}} u^{\bar{\beta}} = F^{\bar{\alpha}}.$$

- Dual-coordinate frame method uses basis vectors of one coordinate system to define components of fields, and a second set of coordinates,  $x^a = \{t, x^i\} = x^a(x^{\bar{a}})$ , to represent these components as functions,  $u^{\bar{\alpha}} = u^{\bar{\alpha}}(x^a)$ .

# Dual-Coordinate-Frame Evolution Method

- Single-coordinate frame method uses the one set of coordinates,  $x^{\bar{a}} = \{\bar{t}, \bar{x}^i\}$ , to define field components,  $u^{\bar{\alpha}} = \{\psi_{\bar{a}\bar{b}}, \Pi_{\bar{a}\bar{b}}, \Phi_{\bar{i}\bar{a}\bar{b}}\}$ , and the same coordinates to determine these components by solving Einstein's equation for  $u^{\bar{\alpha}} = u^{\bar{\alpha}}(x^{\bar{a}})$ :

$$\partial_{\bar{i}} u^{\bar{\alpha}} + A^{\bar{k}\bar{\alpha}}_{\bar{\beta}} \partial_{\bar{k}} u^{\bar{\beta}} = F^{\bar{\alpha}}.$$


- Dual-coordinate frame method uses basis vectors of one coordinate system to define components of fields, and a second set of coordinates,  $x^a = \{t, x^i\} = x^a(x^{\bar{a}})$ , to represent these components as functions,  $u^{\bar{\alpha}} = u^{\bar{\alpha}}(x^a)$ .
- These functions are determined by solving the transformed Einstein equation:

$$\partial_t u^{\bar{\alpha}} + \left[ \frac{\partial x^i}{\partial \bar{t}} \delta^{\bar{\alpha}}_{\bar{\beta}} + \frac{\partial x^i}{\partial x^{\bar{k}}} A^{\bar{k}\bar{\alpha}}_{\bar{\beta}} \right] \partial_i u^{\bar{\beta}} = F^{\bar{\alpha}}.$$


# Testing Dual-Coordinate-Frame Evolutions

- Single-frame evolutions of Schwarzschild in rotating coordinates are unstable, while dual-frame evolutions are stable:

## Dual Frame Evolution



## Single Frame Evolution



- Dual-frame evolution shown here uses a comoving frame with  $\Omega = 0.2/M$  on a domain with outer radius  $r = 1000M$ .

# Horizon Tracking Coordinates


- Coordinates must be used that track the motions of the holes.
- The coordinate transformation from inertial coordinates,  $(\bar{x}, \bar{y}, \bar{z})$ , to co-moving coordinates  $(x, y, z)$ ,

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = e^{a(\bar{t})} \begin{pmatrix} \cos \varphi(\bar{t}) & -\sin \varphi(\bar{t}) & 0 \\ \sin \varphi(\bar{t}) & \cos \varphi(\bar{t}) & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \bar{x} \\ \bar{y} \\ \bar{z} \end{pmatrix},$$

with  $t = \bar{t}$ , is general enough to keep the holes fixed in co-moving coordinates for suitably chosen functions  $a(\bar{t})$  and  $\varphi(\bar{t})$ .


- Since the motions of the holes are not known *a priori*, the functions  $a(\bar{t})$  and  $\varphi(\bar{t})$  must be chosen dynamically and adaptively as the system evolves.

## Horizon Tracking Coordinates II



- Measure the comoving centers of the holes:  $x_c(t)$  and  $y_c(t)$ , or equivalently  $Q^x(t) = [x_c(t) - x_c(0)]/x_c(0)$  and  $Q^y(t) = y_c(t)/x_c(t)$ .
- Choose the map parameters  $a(t)$  and  $\varphi(t)$  to keep  $Q^x(t)$  and  $Q^y(t)$  small.

## Horizon Tracking Coordinates II




- Measure the comoving centers of the holes:  $x_c(t)$  and  $y_c(t)$ , or equivalently  $Q^x(t) = [x_c(t) - x_c(0)]/x_c(0)$  and  $Q^y(t) = y_c(t)/x_c(t)$ .
- Choose the map parameters  $a(t)$  and  $\varphi(t)$  to keep  $Q^x(t)$  and  $Q^y(t)$  small.
- Changing the map parameters by the small amounts,  $\delta a$  and  $\delta\varphi$ , results in associated small changes in  $\delta Q^x$  and  $\delta Q^y$ :

$$\delta Q^x = -\delta a, \quad \delta Q^y = -\delta\varphi.$$



## Horizon Tracking Coordinates II



- Measure the comoving centers of the holes:  $x_c(t)$  and  $y_c(t)$ , or equivalently  $Q^x(t) = [x_c(t) - x_c(0)]/x_c(0)$  and  $Q^y(t) = y_c(t)/x_c(t)$ .
- Choose the map parameters  $a(t)$  and  $\varphi(t)$  to keep  $Q^x(t)$  and  $Q^y(t)$  small.
- Changing the map parameters by the small amounts,  $\delta a$  and  $\delta\varphi$ , results in associated small changes in  $\delta Q^x$  and  $\delta Q^y$ :

$$\delta Q^x = -\delta a, \quad \delta Q^y = -\delta\varphi.$$

- Measure the quantities  $Q^y(t)$ ,  $dQ^y(t)/dt$ ,  $d^2Q^y(t)/dt^2$ , and set

$$\frac{d^3\varphi}{dt^3} = \lambda^3 Q^y + 3\lambda^2 \frac{dQ^y}{dt} + 3\lambda \frac{d^2Q^y}{dt^2} = -\frac{d^3Q^y}{dt^3}.$$

The solutions to this “closed-loop” equation for  $Q^y$  have the form  $Q^y(t) = (At^2 + Bt + C)e^{-\lambda t}$ , so  $Q^y$  always decreases as  $t \rightarrow \infty$ .

## Horizon Tracking Coordinates III

- In practice the coordinate maps are adjusted only at a prescribed set of adjustment times  $t = t_i$ .
- In the time interval  $t_i < t < t_{i+1}$  we set:

$$\begin{aligned} \varphi(t) = & \varphi_i + (t - t_i) \frac{d\varphi_i}{dt} + \frac{(t - t_i)^2}{2} \frac{d^2\varphi_i}{dt^2} \\ & + \frac{(t - t_i)^3}{2} \left( \lambda \frac{d^2 Q_i^y}{dt^2} + \lambda^2 \frac{dQ_i^y}{dt} + \lambda^3 \frac{Q_i^y}{3} \right), \end{aligned}$$

where  $Q^x$ ,  $Q^y$ , and their derivatives are measured at  $t = t_i$ , so these maps satisfy the closed loop equation at  $t = t_i$ .


## Horizon Tracking Coordinates III

- In practice the coordinate maps are adjusted only at a prescribed set of adjustment times  $t = t_i$ .
- In the time interval  $t_i < t < t_{i+1}$  we set:

$$\varphi(t) = \varphi_i + (t - t_i) \frac{d\varphi_i}{dt} + \frac{(t - t_i)^2}{2} \frac{d^2\varphi_i}{dt^2} + \frac{(t - t_i)^3}{2} \left( \lambda \frac{d^2 Q_i^y}{dt^2} + \lambda^2 \frac{dQ_i^y}{dt} + \lambda^3 \frac{Q_i^y}{3} \right),$$


where  $Q^x$ ,  $Q^y$ , and their derivatives are measured at  $t = t_i$ , so these maps satisfy the closed loop equation at  $t = t_i$ .

- **This works!** We are now able to evolve binary black holes using horizon tracking coordinates until just before merger.




## Evolving Binary Black Hole Spacetimes

- We can now evolve binary black hole spacetimes with excellent accuracy and computational efficiency through many orbits.





Head-on Merger Movie



Lapse- $\Psi_4$  Movie


## Evolving Binary Black Hole Spacetimes II

- Gravitational waveform and frequency evolution for the equal mass non-spinning BBH evolution.



# Evolving Binary Black Hole Spacetimes III

- Preliminary error estimates for the gravitational wave phase for the 15 orbit evolution.



# Evolving Binary Black Hole Spacetimes IV

- Preliminary comparisons of the gravitational wave phase for the 15 orbit evolution with various PN order predicted waveforms.

